



U.S. Government Exercises Option to Purchase Additional RAPIVAB® (peramivir injection) from BioCryst for Delivery to Strategic National Stockpile

September 1, 2021

RESEARCH TRIANGLE PARK, N.C., Sept. 01, 2021 (GLOBE NEWSWIRE) -- [BioCryst Pharmaceuticals, Inc.](https://www.biocryst.com) (Nasdaq: BCRX) today announced that the U.S. Department of Health and Human Services (HHS) has exercised its option to purchase an additional 10,000 doses of BioCryst's antiviral influenza therapy, RAPIVAB® (peramivir injection), for approximately \$7 million.

The RAPIVAB purchase by the HHS Office of the Assistant Secretary for Preparedness and Response will supply the Strategic National Stockpile (SNS), the nation's largest supply of life-saving pharmaceuticals and medical supplies for use in a public health emergency.

"The Strategic National Stockpile is an important line of defense in our efforts to ensure availability of critical medical assets to protect the health of Americans in the event of a public health emergency. We are pleased to provide additional doses of RAPIVAB to the SNS as we enter another influenza season of unpredictable severity," said Dr. William Sheridan, chief medical officer of BioCryst.

The order is part of a \$34.7 million contract (Contract No. 75D301-18-C-02984) the Centers for Disease Control and Prevention awarded in 2018 for the procurement of up to 50,000 doses of RAPIVAB over a five-year period for the SNS. With the fulfillment of this new order, BioCryst will have delivered 40,000 doses under the contract.

About RAPIVAB® (peramivir injection)

RAPIVAB® (peramivir injection) is approved in the United States for the treatment of acute uncomplicated influenza in patients six months and older who have been symptomatic for no more than two days. It is administered via an intravenous infusion for a minimum of 15 minutes at recommended doses of 600 mg/kg for adults and adolescents and 12 mg/kg for pediatric patients ages six months to 12 years. Efficacy of RAPIVAB is based on clinical trials of naturally occurring influenza in which the predominant influenza infections were influenza A virus and a limited number of patients infected with influenza B virus. Visit <http://www.rapivab.com> to learn more.

U.S. Indication and Important Safety Information

Indication

RAPIVAB is indicated for the treatment of acute uncomplicated influenza in patients 6 months and older who have been symptomatic for no more than 2 days.

Limitations of Use

- Efficacy of RAPIVAB is based on clinical trials of naturally occurring influenza in which the predominant influenza infections were influenza A virus; a limited number of subjects infected with influenza B virus were enrolled.
- Influenza viruses change over time. Emergence of resistance substitutions could decrease drug effectiveness. Other factors (for example, changes in viral virulence) might also diminish clinical benefit of antiviral drugs. Prescribers should consider available information on influenza drug susceptibility patterns and treatment effects when deciding whether to use RAPIVAB.
- The efficacy of RAPIVAB could not be established in patients with serious influenza requiring hospitalization.

Contraindications

RAPIVAB is contraindicated in patients with known serious hypersensitivity or anaphylaxis to peramivir or any component of the product. Severe allergic reactions have included anaphylaxis, erythema multiforme and Stevens-Johnson Syndrome.

Warnings and Precautions

- Cases of anaphylaxis and serious skin/hypersensitivity reactions such as Stevens-Johnson syndrome and erythema multiforme have occurred with RAPIVAB. Discontinue RAPIVAB and initiate appropriate treatment if anaphylaxis or serious skin reaction occurs or is suspected.
- Neuropsychiatric events: Patients with influenza may be at an increased risk of hallucinations, delirium and abnormal behavior early in their illness. Monitor for signs of abnormal behavior.
- Serious bacterial infections may begin with influenza-like symptoms or may coexist with or occur as complications during the course of influenza. RAPIVAB has not been shown to prevent such complications.

Adverse Reactions

The most common adverse reaction in adults (18 years of age and older) was diarrhea (8% RAPIVAB vs 7% placebo). Lab abnormalities (incidence $\geq 2\%$) occurring more commonly with RAPIVAB than placebo were elevated ALT > 2.5 times the upper limit of normal (3% vs 2%), elevated serum glucose >160 mg/dL (5% vs 3%), elevated CPK at least 6 times the upper limit of normal (4% vs 2%), and neutrophils $<1.0 \times 10^9/L$ (8% vs 6%). In a subset of subjects with serious influenza requiring hospitalization treated with RAPIVAB 600 mg as monotherapy (N=101), the following adverse

reactions were also reported more frequently with RAPIVAB as compared to placebo: constipation (4% versus 2%), insomnia (3% versus 0%), AST increased (3% versus 2%), and hypertension (2% versus 0%).

The safety profile of RAPIVAB in subjects 6 months to 17 years of age was generally similar to that observed in adults. The only adverse reaction reported in pediatric subjects treated with RAPIVAB (occurring in $\geq 2\%$ of subjects) and not reported in adults was vomiting (3% versus 9% for oseltamivir). The only clinically significant laboratory abnormality (DAIDS Grade 2) occurring in $\geq 2\%$ of pediatric subjects treated with RAPIVAB (and not previously reported in adults) was proteinuria by dipstick analysis (3% versus 0% for oseltamivir).

Concurrent Use With Live Attenuated Influenza Vaccine

Antiviral drugs may inhibit viral replication of a live attenuated influenza vaccine (LAIV) and thus may reduce vaccine efficacy. The concurrent use of RAPIVAB with LAIV intranasal has not been evaluated. Avoid use of LAIV within 2 weeks before or 48 hours after administration of RAPIVAB, unless medically indicated.

Please see [full prescribing information](#) for RAPIVAB.

You are encouraged to report negative side effects of prescription drugs to the FDA. To report suspected adverse reactions, contact BioCryst Pharmaceuticals at 1-833-633-2279 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

About BioCryst Pharmaceuticals

BioCryst Pharmaceuticals discovers novel, oral, small-molecule medicines that treat rare diseases in which significant unmet medical needs exist and an enzyme plays a key role in the biological pathway of the disease. Oral, once-daily ORLADEYO[®] (berotralstat) is approved in the United States, the European Union, Japan and the United Kingdom. BioCryst has several ongoing development programs including BCX9930, an oral Factor D inhibitor for the treatment of complement-mediated diseases, BCX9250, an ALK-2 inhibitor for the treatment of fibrodysplasia ossificans progressiva, and galidesivir, a potential treatment for Marburg virus disease and Yellow Fever. RAPIVAB[®] (peramivir injection) has received regulatory approval in the U.S., Canada, Australia, Japan, Taiwan and Korea. Post-marketing commitments for RAPIVAB are ongoing. For more information, please visit the company's website at www.biocryst.com.

Forward-Looking Statements

This press release contains forward-looking statements, including statements regarding sales of RAPIVAB. These statements involve known and unknown risks, uncertainties and other factors which may cause BioCryst's actual sales to be materially different from those expressed or implied by the forward-looking statements. These statements reflect our current views with respect to future events and are based on assumptions and are subject to risks and uncertainties. Given these uncertainties, you should not place undue reliance on these forward-looking statements. Some of the factors that could affect the forward-looking statements contained herein include: the U.S. government may purchase smaller quantities of RAPIVAB than currently anticipated, or none at all; BioCryst relies on third-party manufacturers to manufacture RAPIVAB in a timely manner and in accordance with applicable governmental regulations, and any failure of such third-party manufacturers to perform their obligations could impact BioCryst's ability to supply RAPIVAB pursuant to the government contract; government contracts contain certain terms and conditions, including termination provisions, that subject BioCryst to additional risks; and the ongoing COVID-19 pandemic, which could create challenges in all aspects of BioCryst's business, including without limitation delays, stoppages, difficulties, and increased expenses with respect to BioCryst's and its partners' supply chains, negatively impact BioCryst's ability to access the capital or credit markets to finance its operations, or have the effect of heightening the other risks described herein or in the documents BioCryst files periodically with the Securities and Exchange Commission. Please refer to the documents BioCryst files periodically with the Securities and Exchange Commission, specifically BioCryst's most recent Annual Report on Form 10-K, Quarterly Reports on Form 10-Q, and Current Reports on Form 8-K, all of which identify important factors that could cause the actual results to differ materially from those contained in BioCryst's forward-looking statements.

BCRXW

Investors:

John Bluth
+1 919 859 7910
jbluth@biocryst.com

Media:

Catherine Collier Kyroulis
+1 917 886 5586
ckyroulis@biocryst.com